

1-Alkyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane Salts: a Novel Family of Electrophilic Fluorinating Agents¹

R. Eric Banks,^{*a} Suad N. Mohialdin-Khaffaf,^a G. Sankar Lal,^b Iqbal Sharif^a and Robert G. Syvret^b

^a Chemistry Department, The University of Manchester Institute of Science and Technology, Manchester M60 1QD, UK

^b Corporate Science and Technology Center, Air Products and Chemicals Inc., Allentown, PA 18195-1501, USA

Members of a new series of solid, easily handled, storable, transportable, commercially viable, site-selective electrophilic fluorinating agents of the ⁺N-F class (tradenamed Selectfluor reagents) have been synthesized *via* direct fluorination of monoquaternary salts of 1,4-diazabicyclo[2.2.2]octane.


Unremitting interest worldwide in site-selective fluorination (including ¹⁸F placement) of biologically-active molecules^{2,3} has heightened demand for more generally acceptable (less aggressive, non-explosive, less toxic, inexpensive) electrophilic fluorinating agents⁴ than perchloryl fluoride,⁵ O-F reagents (CF₃OF, CsSO₄F),⁴⁻⁶ xenon difluoride⁷ or fluorine itself.⁸ Of late, the search for an ideal 'F⁺' transfer agent has centred on N-F compounds,⁹ with the result that *N*-alkyl-*N*-fluoro-*p*-toluenesulfonamides (Barnette reagents), *N*-fluoroquinuclidinium fluoride and trifluoromethanesulfonate (triflate, TfO⁻) (Banks reagents) **1** and certain *N*-fluoropyridinium salts (Umemoto reagents), including a modification of the prototypical tetrafluoroborate,¹⁰ are now commercially available.

Direct fluorination of 1,4-diazabicyclo[2.2.2]octane and its monoquaternary salts **2** has been studied with the objective of producing electrophilic fluorinating agents that are just as easy and safe to use as *N*-fluoroquinuclidinium salts **1**¹¹ but more powerful. This simple strategy of incorporating a second quaternized bridgehead nitrogen into the ring system proved extremely successful, for not only did it produce the desired effect, but also it enabled the fluorinating power to be 'tuned' through variation in the electronegativity of the quaternizing group R.


Attempts to prepare 1,4-difluoro-1,4-diazoniabicyclo[2.2.2]octane difluoride **3** (R = F; X⁻ = F⁻) or ditriflate **3** (R = F; X⁻ = TfO⁻) *via* passage, respectively, of neat fluorine at approximately 20 mmHg pressure into cold (-78 °C) trichlorofluoromethane containing 1,4-diazabicyclo[2.2.2]octane,¹² or fluorine-nitrogen blends into cold (-20 to -40 °C) solutions of the same substrate in acetonitrile containing lithium triflate at atmospheric pressure, proved unsatisfactory [cf. ref. 11 the successful syntheses of **1** (X⁻ = F⁻ or TfO⁻)]. However, fluorination of monoquaternary salts **2** (R = Me, CH₂Cl, or CF₃CH₂; and X⁻ = TfO⁻ or BF₄⁻, for example) in the presence of lithium triflate or sodium tetrafluoroborate, as appropriate, proceeded smoothly in cold acetonitrile to give high yields (87-95% after purification) of the corresponding 1-alkyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane salts **3** (R = Me, CH₂Cl or CH₂CF₃; X⁻ = TfO⁻ or BF₄⁻). The structures of these new fluoroammonium salts were established by

elemental (C, H, F, N) and spectroscopic analyses [notably ¹H, ¹³C, ¹⁹F; δ⁺NF (CF₃CO₂H) values (D₂O solutions) fall in the range from +124 to +127.5 (cf. ref. 11a, *N*-fluoroquinuclidinium salts of from +133.5 to +135.5)], and X-ray crystallography [⁺N-F distance in **3** (R = CH₂Cl; X⁻ = BF₄⁻) = 1.37(2) Å;¹³ cf. ref. 13 **1** (X⁻ = TfO⁻) 1.406 Å]; they are white, freeflowing, virtually non-hygroscopic, crystalline, high-melting solids [e.g. **3** (R = CH₂Cl; X⁻ = BF₄⁻) 190 °C; **3** (R = Me; X⁻ = TfO⁻) 220-221 °C (decomp.)] that are soluble in polar solvents [H₂O, Me, MeCN and dimethylformamide (Me₂SO attacks them)], liberate iodine rapidly from acidified (HCl) aqueous acetonic potassium iodide, hence they can be assayed (1 mol ⁺NF ≡ 1 I₂),¹¹ oxidize bromide ion to bromine at room temperature, and, though stable towards cold aqueous mineral acids, are decomposed by cold aqueous sodium hydroxide. Conventional methodology provides easy access to the monoquaternary salts **2** used as precursors of the Selectfluor reagents **3** {e.g. 1,4-diazabicyclo[2.2.2]octane + MeCl + NaBF₄ (or LiOTf) in MeCN at 20 °C → **2**, R = Me, X⁻ = BF₄⁻ (or TfO⁻), in 98% (or 86%) yield, after purification}.

1-Alkyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane salts **3** smoothly and selectively fluorinate a variety of substrates with



Scheme 1 Reagents and conditions: i, testosterone enol diacetate, MeCN, 15 min at 25 °C (R = CH₂Cl, X⁻ = BF₄⁻); ii, androsterone enol diacetate, MeCN, 2 h at 25 °C (R = CH₂Cl, X⁻ = BF₄⁻); iii, 1-morpholinocyclohex-1-ene, CH₂Cl₂, 20 °C (R = CH₂CF₃, X⁻ = TfO⁻); iv, PhMgBr, 25 °C, 2 h, Et₂O (R = CH₂Cl, X⁻ = TfO⁻); v, PhNHCOMe, refluxing MeCN, 15 min (R = CH₂Cl, X⁻ = BF₄⁻); vi, sodio derivative of diethyl phenylmalonate, tetrahydrofuran-dimethylformamide (2:1), 25 °C, 30 min (R = Me, X⁻ = TfO⁻). Reactant ratios = 1:1 molar. Yields refer to isolated products.



overt or masked carbanionic character, as exemplified in the fluorination reactions of Scheme 1. Steroids, in the form of the appropriate enol acetates are rapidly and selectively fluorinated at room temperature giving the 6- or 16-fluoro compounds in high yields, the latter also with very high stereoselectivity ($\alpha : \beta = 95 : 5$). In this application they are far more effective than the other available ^+N-F reagents.^{4,10,11} Highly-stabilized carbanions such as the sodium salt of diethyl phenylmalonate give fluoro derivatives in high yields, but the reagents appear to be less effective with more reactive carbanion systems such as ketone-derived metal enolates. Grignard reagents are smoothly converted to the corresponding monofluoro compounds.

Within the Selectfluor reagent series there is greater ease of 'F⁺' transfer as the electronegativity of the quaternary group R increases. For example, with **3** where R = CH₂CF₃ and X⁻ = TfO⁻, in wet acetonitrile there is 72% conversion of anisole to an approximately 1:1 mixture of *o*- and *p*-fluoroanisole in 5 h at 40 °C. With the *N*-chloromethyl (**3**: R = CH₂Cl; X⁻ = TfO⁻) and *N*-methyl (**3**: R = Me; X⁻ = TfO⁻) analogues, the reaction proceeds to the same extent under otherwise identical conditions in 6 and 13 h, respectively. In consonance with the stratagem on which the discovery of reagents of type **3** was based, *N*-fluoroquinuclidinium salts **1** have been found to be noticeably less reactive than salts of type **3** towards many types of substrate. Thus, fluorination of anisole at 40 °C in wet acetonitrile is imperceptible (gas chromatographic analysis) even after 24 h.

The Selectfluor salts comprise a highly reactive, yet remarkably stable, family of selective electrophilic fluorinating agents. Investigations aimed at uncovering further applications of these new reagents are in progress.

We are indebted to Dr Guido P. Pez (Air Products and Chemicals Inc.) for helpful discussions, and Air Products and Chemicals Inc., for financing the work at UMIST.

Received, 29th January 1992; Com. 2/00500J

References

- 1 Oral presentation at the 13th International Symposium on Fluorine Chemistry (see R. E. Banks and G. S. Lal, abstracts D20 and D21, *J. Fluorine Chem.*, 1991, **54**, 207). The new ^+N-F compounds described here are members of Air Products and Chemicals, Selectfluor family of fluorinating agents; information regarding availability and applications is available from Mr T. A. Shenoy, Air Products and Chemicals Inc., 7201 Hamilton Boulevard, Allentown, PA 18195-1501, USA.
- 2 J. T. Welch and S. Eswarakrishnan, *Fluorine in Bioorganic Chemistry*, John Wiley, New York, 1991.
- 3 M. R. Kilbourn, *Fluorine-18 Labeling of Radiopharmaceuticals*, National Academy Press, Washington DC, 1990.
- 4 For a recent review of electrophilic fluorination, see G. G. Furin in *New Fluorinating Agents in Organic Syntheses*, ed. L. German and S. Zemskov, Springer-Verlag, Berlin, 1989, pp. 34-68; note that extensive lists of primary literature appear in refs. 9 and 11a.
- 5 C. M. Sharts and W. A. Sheppard, *Org. React.*, 1974, **21**, 225.
- 6 F. M. Mukametshin, ref. 4, pp. 59-116.
- 7 V. V. Bardin and Yu. L. Yagupolskii, ref. 4, pp. 1-34.
- 8 S. T. Purrington, B. S. Kagen and T. B. Patrick, *Chem. Rev.*, 1986, **86**, 997.
- 9 V. Murtagh, *Perform. Chem.*, 1991, **6** (4), 35; for extensive, up-to-date lists of primary references, see also R. E. Banks, V. Murtagh and E. Tsiliopoulos, *J. Fluorine Chem.*, 1991, **52**, 389, and G. Resnati and D. D. DesMarteau, *J. Org. Chem.*, 1991, **56**, 4925.
- 10 M. Van Der Puy, D. Nalewajek, G. A. Shia and W. J. Wagner, US Pat. 4,935,519/1990; A. J. Poss, M. Van Der Puy, D. Nalewajek, G. A. Shia, W. J. Wagner and R. L. Frenette, *J. Org. Chem.*, 1991, **56**, 5962. Note, however, that our analysis (including an X-ray study) of a commercial sample of so-called *N*-fluoropyridinium pyridine heptafluorodiborate indicates that the material is *N*-fluoropyridinium pyridinium tetrafluoroborate trifluorohydroxyborate (R. E. Banks, M. K. Besheesh and R. Pritchard, in preparation).
- 11 (a) R. E. Banks, R. A. Du Boisson, W. D. Morton and E. Tsiliopoulos, *J. Chem. Soc., Perkin Trans. 1*, 1988, 2805; (b) R. E. Banks and I. Sharif, *J. Fluorine Chem.*, 1988, **41**, 297; (c) *J. Fluorine Chem.*, 1992, **55**, 207.
- 12 R. E. Banks and E. Tsiliopoulos, unpublished results; E. Tsiliopoulos, PhD Thesis, University of Manchester, 1986.
- 13 R. E. Banks, R. G. Pritchard and I. Sharif, in preparation.
- 14 Thermal analysis revealed that **3** (R = CH₂Cl; X⁻ = BF₄⁻) decomposes exothermally on heating, beginning at 190 °C.